HEDIS[®] Measure Reference Guide



Medicaid NCQA Technical Specifications 2025

What is HEDIS[®]?

HEDIS (Healthcare Effectiveness Data and Information Set) is a set of standardized performance measures developed by the National Committee of Quality Assurance (NCQA) to objectively measure, report, and compare quality across health plans. NCQA develops HEDIS measures through a committee represented by purchasers, consumers, health plans, health care providers, and policy makers.

What are HEDIS scores used for?

As state and federal governments move toward a quality-driven healthcare industry, HEDIS rates are becoming more important for both health plans and individual providers. State purchasers of healthcare use aggregated HEDIS rates to evaluate health insurance companies; efforts to improve preventative care health outreach for members.

How are HEDIS rates calculated?

HEDIS rates can be calculated in two ways: administrative date or hybrid data. Administrative data consists of claim or encounter data submitted to the health plan. Hybrid data consists of both administrative data and a sample of medical record data. Hybrid data requires review of a random sample of member medical records to abstract data for services rendered, but were not reported to the health plan through claims or encounter data. Accurate and timely claim submission/encounter data reduces the need for medical record review. If services are not billed or not billed accurately, they are not included in the calculation.

How can provider HEDIS scores be improved using coding, billing and supplemental data?

- Submit claim/encounter data for each and every service rendered
- Make sure that chart documentation reflects all services billed
- Bill (or report by encounter submission) for all delivered services, regardless of contract status
- Ensure that all claim/encounter data is submitted in an accurate and timely manner
- Consider including CPT II codes to provide additional details and reduce medical record requests
 - Work with the CountyCare Quality Team to learn about submitting supplemental data year-round.
 You can contact us at CountyCarepophealth@cookcountyhhs.org.

Who to outreach for more information about HEDIS?

Contact the Population Health and Performance Improvement Department by email at **countycarepophealth@cookcountyhhs.org** or reach out to your assigned Provider Relations representative. You can also contact Provider Services by calling 312-864-8200.

The guide will serve as a helpful reference tool and is not intended to replace professional coding standards or billing practices. Measures and codes in the HEDIS Measure Reference Guide are not all-inclusive and can be changed, deleted or removed at any time. Measures are derived from the NCQA HEDIS Measurement Year 2025 Technical Specifications.

HEDIS® is a registered trademark of the National Committee for Quality Assurance (NCQA).



Table of Contents

Definitions	. 3
Timeline	. 3
Telehealth & In-Person Visits	. 4
Preventive Care . Adults' Access to Preventive/Ambulatory Health Services (AAP) . Adult Immunization Status (AIS-E) . Colorectal Cancer Screening (COL-E) . Breast Cancer Screening (BCS-E) . Cervical Cancer Screening (CCS-E) . Chlamydia Screening (CHL) .	. 5 . 6 . 7 . 8 . 9
Keeping Kids Healthy Child and Adolescent Well-Care Visits (WCV). Childhood Immunization Status (CIS-E) Immunizations for Adolescents (IMA-E) Lead Screening in Children (LSC) Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC) Well-Child Visits in the First 30 Months of Life (W30)	10 11 12 12 13
Pregnant Members Prenatal and Postpartum Care (PPC) Prenatal Immunization Status (PRS-E)	14
Living with Chronic Conditions Cardiac Rehabilitation (CRE) Controlling Blood Pressure (CBP) Persistence of Beta-Blocker Treatment After a Heart Attack (PBH). Pharmacotherapy Management of COPD Exacerbation (PCE) Plan All – Cause Readmission (PCR)	16 17 17 18
Diabetes Management. Blood Pressure Control for Patients with Diabetes (BPD)*. Eye Exam for Patients With Diabetes (EED)* Glycemic Status Assessment for Patients with Diabetes (GSD) Kidney Health Evaluation for Patients with Diabetes (KED). Statin Therapy for Patients with Diabetes (SPD)	20 20 20 22
Behavioral Health. Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC) Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD) Diabetes Screening for People with Schizophrenia or	24
Bipolar Disorder who are Using Antipsychotic Medications (SSD) Follow-Up After Emergency Department Visit for Substance Use Disorder or Dependence (FUA) Follow-Up After Emergency Department Visit for Mental Illness (FUM) Substance Use Disorder (FUI) Follow-Up After Hospitalization for Mental Illness (FUH) Follow-Up Care for Children Prescribed ADHD Medication (ADD-E) Initiation & Engagement of Alcohol and Other Drug Abuse or Independence Treatment (IET). Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM) Pharmacotherapy for Opioid Use Disorder (POD)	25 27 30 32 33 33



Definitions

- Measurement Year (MY) 2025 The 12-month timeframe between which a service was rendered – generally January 1 through December 31. Data collected from this timeframe is reported during the reporting year (2026).
- Reporting Year The timeframe when data is collected and reported. The service dates are from the measurement year, which is usually the year prior. In some cases, the service dates may go back more than one year. For example: The 2026 reporting year would include data from services rendered during the measurement year, which would be 2025 and/or any time prior.

Results from the 2026 reporting year would likely be released in June 2026.

- 3. **Denominator** The number of members who qualify for the measure criteria, based on NCQA technical specifications.
- Numerator The number of members who meet compliance criteria based on NCQA technical specifications for appropriate care, treatment or service.
- Reporting Methodology The method a health plan uses to collect and report data to calculate measures. Three methods are used: Administrative, Hybrid, and ECDS.
 - ECDS (Electronic Clinical Data Systems) a reporting method used to collect and report structured electronic clinical quality data.

ECDS Measures required for MY2025:

- Breast Cancer Screening (BCS-E)
- Cervical Cancer Screening (CCS-E)
- Childhood Immunization Status (CIS-E)
- Immunizations for Adolescents (IMA-E)
- Colorectal Cancer Screening (COL-E)
- Medical Record Data The information taken directly from a member's medical record to validate services rendered that weren't captured through medical or pharmacy claims, encounters, or supplemental data.
- Required Exclusions Members are excluded from a measure denominator based on their diagnosis and/or procedure captured in claim/encounter data. A determination is made after the claim is processed within certified HEDIS software while the measure denominator is being created. (Applicable to all measures)

For example:

- Members with a claim for hospice services during the measurement year will be excluded from all applicable measures.
- Members who die any time during the measurement year will be excluded from all applicable measures.
- Optional Exclusion Members are excluded from a measure denominator manually using certified HEDIS software during the hybrid review process, also known as medical records review.

Timeline

Annual HEDIS Timeline

January to early May	June	September/October
Performance Improvement and Popu- lation Health department staff collect and request and review medical records from previous year.	HEDIS results are certified and reported to NCQA.	NCQA release Quality Compass results nationwide.

CountyCare

Telehealth & In-Person Visits

The following measures can be met via an in-person, telephone, telehealth visit and/or e-visit (virtual check-in):

HEDIS Measure	In-person Visit	Telephone Visit	Telehealth Visit	E-Visit
Adults' Access to Preventive/Ambulatory Health Services (AAP)	1	1		
Follow-Up Care for Children Prescribed ADHD Medication (ADD – Continuation)	1	1	5	
Blood Pressure for Patients with Diabetes (BPD)	1	1	1	
Controlling Blood Pressure (CBP)	1	1	1	
Follow Up After Emergency Department Visit for Alcohol and Other Drug Abuse or Dependence (FUA)	1		\$	
Follow Up After Hospitalization for Mental Illness (FUH)	1	1	1	
Follow Up After High-Intensity Care for Substance Use Disorder (FUI)	1	1	1	
Initiation & Engagement of Alcohol and Other Drug Abuse or Independence Treatment (IET)	1	1	1	
Transitions of Care (TRC)	1	1	1	
Prenatal and Postpartum Care (PPC)	1	1	✓	<i>✓</i>

Telehealth visit: Real-time interactive audio AND video telecommunication.

E-visits (also referred to as virtual check-ins): It is not real-time, but requires two-way interaction between the member and the provider. Examples include: patient portal, secure text messaging or email.



Preventive Care

Adults' Access to Preventive/Ambulatory Health Services (AAP)

The percentage of adults who had an ambulatory or preventive care visit during the measurement year.

- Member Population: Member ages 20 and older as of December 31 of the measurement year who had one or more ambulatory or preventive care visit during the measurement.
- CountyCare reports three age stratifications and a total rate:
 - 20-44 years old
 - 45-64 years old
 - 65 and older
 - Total

Description	СРТ	HCPCS	ICD-10
Ambulatory visits	99242–99245, 99304–99310, 99347–99350, 99381–99387, 99391–99397, 99401–99404, 99411, 99412, 99429, 99483, 92002, 92004, 92012, 92014, 99202–99205, 99211–99215, 99318, 99324–99328, 99334– 99337, 98969, 99444, 99483, 99315, 99316, 99341, 99345	G0402, G0438, G0439, G0463, T1015, S0620, S0621	Z00.00, Z00.01, Z00.121, Z00.129, Z00.3, Z00.5, Z00.8, Z02.0–Z02.6, Z02.71, Z02.79, Z02.81–Z02.84, Z02.89, Z02.9, Z76.1, Z76.2
Telephone visits	98966–98968, 99441–99443		
E-visit or virtual check-ins	98970–98972, 98980, 98981, 98981, 99421–99423, 99421– 99243, 99457, 99458	G0071, G2010, G2012, G2250–G2252	



Adult Immunization Status (AIS-E)

The percentage of members 19 and older who are up to date on recommended vaccines for Influenza, Tdap, Hepatitis B, Zoster, and Pneumococcal.

- Required exclusions apply
- Members who had a contraindication for a specific vaccine

Description	СРТ	HCPCS
Adult Influenza	90653, 90662, 90694, 90756, 90674, 90689, 90688, 90686, 90630, 90682, 90661, 90658, 90656, 90654, 90673, 90672, 90660	
Tdap	90714, 90715	
Нер В	90743, 90739, 90759, 90746, 90740, 90747, 90744	
Herpes Zoster	90750	
Pneumococcal	90670, 90671, 90677, 90732	G0009



Colorectal Cancer Screening (COL-E)

*Reported as an ECDS measure

The percentage of members who had appropriate screening for colorectal cancer. Any of the following meet criteria:

- Fecal occult blood test (FOBT) during the MY: guaiac based (gFOBT) / immunochemical FOBT or fecal immunological test (FIT).
- Flexible sigmoidoscopy during the measurement year or the four years prior to the measurement year.
- Colonoscopy during the measurement year or the nine years prior.
- Computerized tomography (CT) colonography during the measurement year or four years prior.

- Fecal immunochemical test (FIT)-DNA (Cologuard®) test during the MY or two years prior.
- Measure population: Medicaid Medicare members ages 45–75 as of December 31 of the MY.

- Required exclusions apply. See page 3.
- Members who have a history of colorectal cancer (cancer of the small intestine does not count).
- Members who had a total colectomy (partial or hemicolectomies do not count).
- Medicare members ages 66 and older enrolled in I-SNP or living long-term in an institution.
- Members age 66 and older with frailty and advanced illness.

Description	СРТ	HCPCS	ICD-10
Colonoscopy	44388–44392, 44394, 44401–44408, 45378–45382, 45384–45386, 45388–45393, 45398	G0105, G0121	
CT colonography	74261–74263		
FIT-DNA test	81528	G0464	
Flexible sigmoidoscopy	45330–45335, 45337–45342, 45346, 45347, 45349–45350	G0104	
Fecal occult blood test (FOBT)	82270, 82274	G0328	
Exclusion: Colorectal cancer			C18.0–C18.9, C19, C20, C21.2, C21.8, C78.5, Z85.038, Z85.048
Exclusion: Total colectomy	44150–44153, 44155–44158, 44210–44212		ODTEOZZ, ODTE4ZZ, ODTE8ZZ, ODTE7ZZ



Breast Cancer Screening (BCS-E)

*Reported as an ECDS measure

The percentage of members who were screened for breast cancer with a mammogram anytime on or between October 1 two years prior to the MY and December 31 of the MY.

- Required exclusions apply. See page 3.
- Members with a history of bilateral mastectomy or both right and left unilateral mastectomies.
- Members who had gender-affirming chest surgery with a diagnosis of gender dysphoria.

Description	СРТ	Modifier	HCPCS	ICD-10
Mammography	77061–77063, 77065–77067			
Exclusion: Bilateral mastectomy				0HTV0ZZ
Exclusion: History of bilateral mastectomy				Z90.13
Exclusion: Unilateral mastectomy with a bilateral modifier	19180, 19200, 19220, 19240, 19303–19307	50		
Exclusion: Unilateral mastectomy with left/right side modifier	19180, 19200, 19220, 19240, 19303–19307	LT, RT		
Exclusion: Left and right unilateral mastectomy				OHTUOZZ, OHTTOZZ
Exclusion: Absence of both right and left breast				Z90.11, Z90.12
Exclusion: Palliative care encounter			G9054, M1017	Z51.5
Exclusion: Gender-affirming chest surgery (CPT 19318) requires CPT 19318 and one ICD-10 code for gender dysphoria)	19318			F64.1, F64.2, F64.8, Z87.890



Cervical Cancer Screening (CCS-E)

*Reported as an ECDS measure

The percentage of members who were screened for cervical cancer with age appropriate cervical cytology and/or high-risk human papillomavirus (hrHPV) testing performed.

Exclusions:

- Required exclusions apply. See page 3.
- History of hysterectomy with no residual cervix.
- Cervical agenesis or acquired absence of cervix.
- Members with sex assigned male at birth.

Description	СРТ	HCPCS	ICD-10
For ages 21–64, a cervical cytology is performed every three years	88141–88143, 88147, 88148, 88150, 88152–88154, 88153, 88164–88167, 88174, 88175	G0123, G0124, G0141, G0143–G0144, G0145, G0147, G0148, P3000, P3001, Q0091	
For ages 30–64, hrHPV testing is performed every five years (can also be part of cotesting with cervical cytology every five years)	87624, 87625	G0476	
Exclusion: Members with a hysterectomy without a residual cervix, <u>or</u> cervical agenesis <u>or</u> acquired absence of the cervix are exempt from this measure	51925, 56308, 57530, 57531, 57540, 57545, 57550, 57555, 57556, 58150, 58152, 58200, 58210, 58240, 58260, 58262, 58263, 58267, 58270, 58275, 58280, 58285, 58290–58294, 58548, 58550, 58552–58554, 58570–58573, 58575, 58951, 58953, 58954, 58956, 59135		Q51.5, Z90.710, Z90.712, OUTCOZZ, OUTC4ZZ, OUTC7ZZ, OUTC8ZZ
Exclusion: Member with sex assigned at birth as male	Sex assigned at birth: LOINC code Of Male: LOINC code LA2-8	e 76689-9	

Chlamydia Screening (CHL)

The percentage of members 16–24 who were recommended for routine chlamydia screening, were identified as sexually active and had one or more chlamydia tests in the MY.

Description	СРТ
Chlamydia tests	87110, 87270, 87320, 87490–87492, 87810

- Required exclusions apply. See page 3.
- Members sex assigned male at birth.



Keeping Kids Healthy

Child and Adolescent Well-Care Visits (WCV)

The percentage of children and adolescents who had one or more comprehensive well-care visits with a PCP or an OB/GYN practitioner during the measurement year. Measure population: Members ages 3–21 as of December 31 of the measurement year.

Description	СРТ	HCPCS	ICD-10
Well-care visit	99382, 99383, 99384, 99385–99392, 99393, 99394, 99395	G0438, G0439, S0302, S0610, S0612, S0613	Z00.00, Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z00.3, Z02.5, Z76.1, Z76.2, Z01.411, Z01.419



Childhood Immunization Status (CIS-E)

*Reported as an ECDS measure

The percentage of children age 2 who receive the required childhood immunization status Combination 10 vaccinations.

Measure population: Children who turn two years old during the measurement year.

Note: Refer to the Illinois Comprehensive Automated Registry Exchange (I-CARE) at I-CARE (illinois.gov) for information on tracking and submitting patient immunization records.

• Combination 10. The percentage of 2-year-old children who had four diphtheria, tetanus and acellular pertussis (DTaP); three polio (IPV); one

measles, mumps and rubella (MMR); three haemophilus influenza type B (HiB); three hepatitis B (HepB); one chicken pox (VZV); four pneumococcal conjugate (PCV); one hepatitis A (HepA); two or three rotavirus (RV); and two influenza (flu) vaccines by their 2nd birthday.

- Required exclusions apply. See page 3.
- Members who had a contraindication for a specific vaccine are excluded from the denominator for all antigen rates and the combination rates.
- Exclusion must be met prior to the child's 2nd birthday.

Description	СРТ	HCPCS	ICD-10
DTaP	90697, 90698, 90700, 90723		
HiB	90644, 90647, 90648, 90697, 90698, 90748		
HepB vaccine or history of hepatitis B illness	90697, 90723, 90740, 90744, 90747, 90748	G0010	B16.0, B16.1, B16.2, B16.9, B17.0, B18.0, B18.1, B19.10, B19.11
IPV	90697, 90698, 90713, 90723		
MMR vaccine or history of measles, mumps or rubella	90707, 90710		B05.0, B05.1, B05.4, B05.81, B05.89, B05.9, B26.0, B26.3, B26.81, B26.85, B26.89, B26.9, B06.00, B06.02, B06.09, B06.81, B06.82, B06.89, B06.9
PCV	90670	G0009	
VZV vaccine or history of varicella zoster illness	90710, 90716		B01.0, B01.11, B01.12, B01.2, B01.81, B01.89, B01.9, B02.0, B02.1, B02.21, B02.24, B02.29, B02.34, B02.39, B02.7, B02.9
HepA vaccine or history of HepA illness	90633		B15.0, B15.9
Flu (one of the two flu vaccines can be a LAIV vaccine administered on the 2nd birthday)	90655, 90657, 90660, 90661, 90672, 90674, 90685–90689, 90756		
RV two-dose schedule	90681		
RV three-dose schedule	90680		



Immunizations for Adolescents (IMA-E)

*Reported as an ECDS measure

The percentage of adolescents who received the required combination 1 and combination 2 vaccinations by their 13th birthday.

Measure population: Members who turn age 13 during the measurement year.

Note: Refer to the Illinois Comprehensive Automated Registry Exchange (I-CARE) at I-CARE (illinois.gov) for information on tracking and submitting patient immunization records.

• Combo 1. The percentage of adolescents age 13 who had at least one dose of meningococcal vaccine and one dose of tetanus, diphtheria toxoids and acellular pertussis (Tdap) vaccine. • Combo 2. The percentage of adolescents age 13 who had at least one dose of meningococcal vaccine, one dose tetanus, diphtheria toxoids and acellular pertussis (Tdap) vaccine, and the complete human papillomavirus (HPV) vaccine series by their 13th birthday.

Exclusions:

- Required exclusions apply. See page 3.
- Members who are in hospice.
- Members who have an anaphylactic reaction to any particular vaccine or its components anytime on or before their 13th birthday.
- Tdap: Members who have encephalopathy with a vaccine adverse-effect code on or before their 13th birthday.

Description	СРТ
Meningococcal serogroups A, C, W, Y vaccine or Meningococcal pentavalent serogroups A, C, W, Y and B (between member's 11th and 13th birthdays)	90619, 90623, 90733, 90734
Tdap vaccine (between member's 10th and 13th birthdays)	90715
2 HPV vaccines (at least 146 days apart on or between the member's 9th and 13th birthdays) OR	90649–90651
3 HPV vaccines (with different dates of service on or between the member's 9th and 13th birthdays)	

Lead Screening in Children (LSC)

The percentage of children age 2 who had one or more capillary or venous lead blood tests for lead poisoning by their 2nd birthday.

Labs and health care providers should report all positive results electronically to the Illinois Department of Public Health's (IDPH's) Illinois Lead Program. Refer to the IDPH website for more information on reporting blood lead levels as required. Measure population: Children should be tested for lead at ages 12 months and 24 months, or when there is no documented lead testing for children up to ages 72 months.

Description	СРТ
Lead test	83655



Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents (WCC)

The percentage of children and adolescents who had an outpatient visit with a PCP or OB/GYN and who had evidence of the following during the measurement year.

- BMI percentile documentation. Medical record documentation must include height, weight and the BMI percentile as a specific value (e.g., 80th percentile) or plotted on an age-growth chart.
- Counseling for Nutrition.* Medical record documentation must include either discussion or counseling of nutrition.

- Counseling for Physical Activity.* Medical record documentation must include either discussion or counseling of physical activity.
- * Services rendered do not require specific settings, a telephone visit, e-visit or virtual check-in meet criteria.

Measure population: Children and adolescents ages 3–17 as of December 31 of the measurement year.

Required exclusions apply. See page 3.

Description	СРТ	HCPCS	ICD-10
BMI percentile documentation			Z68.51–Z68.54
Nutrition counseling	97802–97804	G0270, G0271, G0447, S9449, S9452, S9470	Z71.3
Physical activity counseling		G0447, S9451	Z02.5, Z71.82

Well-Child Visits in the First 30 Months of Life (W30)

The percentage of children who had the required number of comprehensive well-child visits with a PCP during the first 30 months of life.

Two rates are reported for this measure:

- Well-Child Visits in the First 15 Months. Children who turned 15 months old during the measurement year with six or more well-child visits.
- Well-Child Visits for Ages 15–30 Months. Children who turned 30 months old during the measurement year with two or more well-child visits.

Measure population: Medicaid children who turn 15 or 30 months of age during the measurement year.

Description	СРТ	HCPCS	ICD-10
Well-Care Visit	99381, 99382, 99385, 99384, 99383, 99391, 99392–99395, 99461		Z00.00, Z00.01, Z00.110, Z00.111, Z00.121, Z00.129, Z00.2, Z00.3, Z01.411, Z01.419, Z02.5, Z02.84, Z76.1, Z76.2



Pregnant Members

Prenatal and Postpartum Care (PPC)

The percentage of deliveries of live births that received timely perinatal care visits.

Measure population: Members who had deliveries or live birth that occurred between October 8 of the year prior to October 7 of the measurement year. • Timeliness of Prenatal Care: The percentage of deliveries that received a prenatal care visit in the first trimester, on the enrollment start date or within 42 days of enrollment in the organization.

For both rates, services provided during a telephone visit, e-visit, or virtual check-in are eligible for use.

Description	СРТ	CPT-CAT-II	HCPCS	ICD-10
Standalone prenatal visits	99500	0500F, 0501F, 0502F (do not include modifiers 1P, 2P, 3P or 8P)	H1000–H1004	
Prenatal visits with pregnancy- related diagnosis code	99202–99205, 99211– 99215, 99242–99245, 99483		G0463, T1015	
Pregnancy diagnosis				Refer to the current ICD–10 manual for the appropriate pregnancy diagnosis codes.
Prenatal bundle services with date prenatal care initiated	59400, 59425, 59426, 59510, 59610, 59618		H1005	
Telephone visits with a pregnancy- related diagnosis code	98966–98968, 99441–99443			
E-visits or virtual check-ins with pregnancy-related diagnosis code	98970–98972, 98980, 98981, 99421–99423, 99457, 99458		G0071, G2010, G2012, G2250– G2252	



Prenatal and Postpartum Care (PPC) (continued)

• Postpartum Care. The percentage of deliveries that had a postpartum visit on or between 7 and 84 days after delivery.

Description	СРТ	HCPCS	ICD-10
Postpartum visits	57170, 99501, 58300, 59430, 0503F (do not include modifiers 1P, 2P, 3P or 8P)	G0101	
Postpartum care encounter			Z01.411, Z01.419, Z01.42, Z30.430, Z39.1, Z39.2
Cervical cytology	88147, 88148, 88142, 88174, 88143, 88175, 88141, 88164, 88166, 88167, 88165, 88150, 88152, 88153	G0147, G0148, G0141, G0124, G0123, G0143, G0145, G0144, P3000, P3001, Q0091	
Postpartum bundled services with date postpartum care initiated	59622, 59515, 59510, 59618, 59400, 59610, 59410, 59614		
Telephone or telehealth visits	98966–98968, 99441–99443		
E-visits or virtual check-ins	98970–98972, 98980, 98981, 99421–99423, 99457, 99458	G0071, G2010, G2012, G2250–G2252	

Prenatal Immunization Status (PRS-E)

The percentage of deliveries in which members received a flu and Tdap vaccine (tetanus, diphtheria toxoids and acellular pertussis).

Measure population: Members who had deliveries or live birth that occurred in the measurement year (Jan 1 – Dec 31).

Description	СРТ
Tdap Vaccine	90715
Adult Influenza	90688, 90686, 90630, 90682



Living with Chronic Conditions

Cardiac Rehabilitation (CRE)

The percentage of members who completed rehabilitation sessions following a severe or acute qualifying cardiac event.

Four rates are reported:

- Initiation. The percentage of members who attended two or more cardiac rehabilitation sessions within **30 days**.
- Engagement 1. The percentage of members who attended 12 or more cardiac rehabilitation sessions within 90 days.
- Engagement 2. The percentage of members who attended 24 or more cardiac rehabilitation sessions within 180 days.
- Achievement. The percentage of members who attended 36 or more cardiac rehabilitation sessions within 180 days.

Measure population: Members ages 18 and older as of the qualifying cardiac event that occurred on July 1 of the year prior to June 30 of the measurement year. The date of the most recent cardiac event is used.

- Required exclusions apply. See page 3.
- Members who had additional discharges due to cardiac event within 180 days from qualifying event.
- Members who are in hospice or receiving palliative care during measurement year.
- Medicare members ages 66 and older as of December 31 of the measurement year who were enrolled in I-SNP or living long term in an institution.
- Members ages 66–80 as of December 31 of the measurement year with frailty and advanced illness.
- Members ages 81 and older as of December 31 with frailty.

Description	СРТ	HCPCS	ICD-10
Cardiac rehabilitation	93797, 93798	G0422, G0423, S9472	
Myocardial infarction (MI)			I21.01, I21.02, I21.09, I21.11, I21.19, I21.21, I21.29, I21.3, I21.4, I21.9, I21.A1, I21. A9, I22.0–I22.2, I22.8, I22.9, I23.0–I23.8, I25.2
Coronary artery bypass grafting (CABG)	33510–33519, 33521–33523, 33530, 33533–33536	S2205–S2209	
Heart transplant	33927, 33928, 33935, 33945		
Heart valve repair or replacement	33361–33369, 33390, 33391, 33404–33406, 33410–33420, 33422, 33425–33427, 33430, 33440, 33460, 33463–33465, 33468, 33470, 33471, 33474, 33475, 33476, 33477, 33478		
Percutaneous coronary intervention (PCI)	92920, 92924, 92928, 92933, 92937, 92941, 92943	C9600, C9602, C9604, C9606, C9607	
Palliative care encounter		G9054, M1017	Z51.5



Controlling Blood Pressure (CBP)

The percentage of members with a diagnosis of hypertension (HTN) whose blood pressure was adequately controlled (< 140/90 mm Hg) during the measurement year. Members had at least two outpatient visits with a diagnosis of HTN on or between January 1 of the year prior to the measurement year and June 30 of the measurement year.

Measure population: Members ages 18–85 with hypertension.

Note: Remote measurements by any digital device are acceptable. Member reported blood pressure documented in the member's medical record are eligible for reporting.

Exclusions:

- Required exclusions apply. See page 3.
- Members ages 66–80 with frailty and advanced illness.
- Members ages 81 and older with 2 indications of frailty.
- Members who have a diagnosis of pregnancy.
- Members who have evidence of (ESRD) end-stage renal disease or had a kidney transplant or total nephrectomy or dialysis.

Description	CPT-CAT-II	ICD-10
Essential hypertension		110
Systolic < 130	3074F	
Systolic 130–139	3075F	
Systolic ≥ 140	3077F	
Diastolic < 80	3078F	
Diastolic 80–89	3079F	
Diastolic ≥ 90	3080F	

Persistence of Beta-Blocker Treatment After a Heart Attack (PBH)

The percentage of members hospitalized and discharged from July 1 of the year prior to the measurement year to June 30 of the measurement year due to acute myocardial infarction (AMI) and who received persistent beta-blocker treatment for six months after discharge. This measure is based on a calculation and there are no codes associated with beta-blocker medications.

Measure population: Members ages 18 and older as of December 31 of the measurement year.

- Required exclusions apply. See page 3.
- Members ages 66–80 as of December 31 of the measurement year with frailty and advanced illness.
- Member ages 81 and older as of December 31 with frailty during the measurement year.

- Members having the following:
 - Asthma
 - Chronic obstructive pulmonary disease (COPD)
 - Obstructive chronic bronchitis
 - Chronic respiratory conditions due to fumes and vapors
 - Hypotension, heart block > 1 degree or sinus bradycardia
 - A medication dispensing event indicative of a history of asthma
 - Intolerance or allergy to beta-blocker therapy

Description	ICD-10
Acute myocardial infraction (AMI)	I21.01, I21.02, I21.09, I21.11, I21.19, I21.21, I21.29, I21.3, I21.4



Pharmacotherapy Management of COPD Exacerbation (PCE)

The percentage of chronic obstructive pulmonary disease (COPD) exacerbations resulting in an acute inpatient discharge or emergency department (ED) visit for the member and had appropriate medications dispensed.

The inpatient discharge or ED visit due to COPD occurred between January 1–November 30 of the measurement year with the following actions:

• Dispensed a systemic corticosteroid (or there was evidence of an active prescription) within 14 days of the event.

Plan All – Cause Readmission (PCR)

The number of acute inpatient stay discharges between January 1 and December 1 during the measurement year that were followed by an unplanned acute readmission within **30 days**. Includes the predicted probability of an acute readmission.

Data for this measure are reported in the following categories:

- Count of Index Hospital Stays (IHS) (denominator)
- Count of Observed 30-Day Readmissions (numerator)
- Count of Expected 30-Day Readmissions

• Dispensed a bronchodilator (or there was evidence of an active prescription) within **30 days** of the event.

Measure population: Members ages 40 or older as of January 1 of the measurement year.

There are no codes for numerator compliance; this is the reason why the list of bronchodilator medications was the only information in previous QRGs.

Required exclusions apply. See page 3.

Measure population:

- Medicaid members ages 18–64 as of January 1 of the measurement year.
- Medicare members ages 18 and older as of January 1 of the measurement year.

Note: A lower rate indicates better performance.

This measure is based on a calculation and there are no codes associated.



Statin Therapy for Patients with Cardiovascular Disease (SPC)

The percentage of members who were identified as having clinical atherosclerotic cardiovascular disease (ASCVD) and who received and adhered to statin therapy.

- Received Statin Therapy. Members who were dispensed at least one high-intensity or moderate-intensity statin medication during the measurement year.
- Statin Adherence 80%. Members who remained on a high-intensity or moderate-intensity statin medication for at least 80% of the treatment period.

Measure population: Males ages 21–75 and females ages 40–75 as of December 31 of the measurement year.

Exclusions:

- Required exclusions apply. See page 3.
- Members diagnosed with myalgia, myositis, myopathy or rhabdomyolysis, or receiving palliative care during the measurement year.
- Members diagnosed with cardiovascular disease, pregnancy, cirrhosis, ESRD or dialysis, in vitro fertilization, or who were dispensed one or more prescriptions for clomiphene during the measurement year and the year prior.
- Members ages 66 and older as of December 31 of the measurement year with frailty and advanced illness.

There are no codes for numerator compliance, just that the member be on a high- or moderate-intensity statin medication during the measurement year.



Diabetes Management

Measure population: Members with diabetes (types 1 and 2) ages 18–75 as of December 31 of the measurement year.

Required exclusions apply. See page 3.

Optional exclusions: Members who had a diagnosis of polycystic ovarian syndrome, gestational diabetes or steroid-induced diabetes during the measurement year.

Blood Pressure Control for Patients with Diabetes (BPD)*

The percentage of members with diabetes who had BP control (< 140/90 mm Hg).

Remote measurements by any digital device are acceptable. Member reported blood pressure documented in the member's medical record are eligible for reporting.

Eye Exam for Patients With Diabetes (EED)*

Administrative reporting method only

The percentage of members with diabetes who had a retinal or dilated eye exam by an eye care professional (optometrist or ophthalmologist) or retinal imaging by a qualified reading center by any provider type.

Glycemic Status Assessment for Patients with Diabetes (GSD)

The percentage of members with diabetes whose most recent HbA1c testing was at the following levels during the measurement year:

- HbA1c control (< 8.0%). The percentage of members with diabetes who had HbA1c control (< 8.0%).
- HbA1c poor control (> 9.0%). The percentage of members with diabetes who had HbA1c poor control (> 9.0%).

Note: The member is not numerator if there is a missing result or if an A1c test was not performed.

Description	CPT-CAT-II
Systolic blood pressure < 130 mmHg	3074F
Systolic blood pressure 130–139 mmHg	3075F
Systolic blood pressure > 140 mmHg	3077F
Diastolic blood pressure < 80 mmHg	3078F
Diastolic blood pressure 80–89 mmHg	3079F
Diastolic ≥ 90 mm Hg	3080F



Diabetes Management (continued)

Description	СРТ	Modifier	CPT-CAT-II	ICD-10	HCPCS
Retinal eye exams	92002, 92004, 92012, 92014, 92018, 92019, 92134, 92201, 92202, 92230, 92235, 92240, 92250, 92260, 99203–99205, 99213–99215, 99242–99245, 98980, 98981				S0620, S0621, S3000
Diabetic retinal screening negative in prior year			3072F		
Diabetes without evidence of complications			E10.9, E11.9, E13.9		
Retinal Imaging	92227, 92228				
Eye exam with interpretation by an ophthalmologist or optometrist documented and reviewed; with evidence of retinopathy			2022F, 2024F, 2026F		
Eye exam with interpretation by an ophthalmologist or optometrist documented and reviewed; without evidence of retinopathy			2023F, 2025F, 2033F		
Unilateral eye enucleation (Unilateral Eye Enucleation Value Set) with a bilateral modifier (Bilateral Modifier Value Set)	65091, 65093, 65101, 65103, 65105, 65110, 65112, 65114				
Unilateral eye enucleation left			08T1XZZ		
Unilateral eye enucleation right			08T0XZZ		
Bilateral Modifier		50			
HbA1c Tests	83036, 83037				
HbA1c test level less than 7.0%			3044F		
HbA1c test level \geq 7.0% and < 8.0%			3051F		
HbA1c test level > 8.0% and < 9.0%			3052F		
HbA1c tests level < 9.0% A lower HbA1c poor control (> 9.0%) rate indicates better performance			3046F		



Kidney Health Evaluation for Patients with Diabetes (KED)

The percentage of members with type 1 and type 2 diabetes who received a kidney health evaluation during the measurement year, with evidence of BOTH of the following:

- An estimated glomerular filtration rate (eGFR)
- Both a quantitative urine albumin lab test and a urine creatinine lab test with service dates four days apart or less

Measure population: Members ages 18–85 with diabetes as of December 31 of the measurement year.

- Required exclusions apply. See page 3.
- Members with ESRD, dialysis or palliative care.
- Medicare members ages 66 and older as of December 31 of the measurement year who were enrolled in I-SNP or living long term in an institution.
- Members ages 66–80 as of December 31 of the measurement year with frailty and advanced illness.
- Members ages 81 and older as of December 31 with frailty.

Description	СРТ	HCPCS	ICD-10CM
eGFR	80047, 80048, 80050, 80053, 80069, 82565		
Quantitative urine albumin lab test	82043		
Urine creatinine lab test	82570		
Exclusion: ESRD			N18.5, N18.6, Z99.2
Exclusion: Dialysis	90935, 90937, 90945, 90947, 90997, 90999, 99512	G0257, S9339	



Statin Therapy for Patients with Diabetes (SPD)

The percentage of adults with diabetes who do not have clinical atherosclerotic cardiovascular disease (ASCVD) who met the following criteria.*

- Received Statin Therapy: Members who were dispensed at least one statin medication of any intensity during the measurement year.*
- Statin Adherence 80%: Members who remained on a statin medication of any intensity for at least 80% of the treatment period.*

Measure population: Members ages 40–75 as of December 31 of the measurement year.

- Required exclusions apply. See page 3.
- Members diagnosed with myalgia, myositis, myopathy or rhabdomyolysis, or receiving palliative care during the measurement year.
- Members diagnosed with cardiovascular disease, pregnancy, cirrhosis, ESRD or dialysis, in vitro fertilization, or was dispensed one or more prescriptions for clomiphene during the measurement year and the year prior.
- Members ages 66 or older as of December 31 of the measurement year with frailty and advanced illness.
- * There are no codes for numerator compliance, just that the member be on a statin medication during the measurement year.



Behavioral Health

Cardiovascular Monitoring for People with Cardiovascular Disease and Schizophrenia (SMC)

The percentage of adults diagnosed with schizophrenia and heart disease who had a cholesterol test during the measurement year. Measure population: Members ages 18–64.

Required exclusions apply. See page 3.

Decription	СРТ	CPT-CAT-II
LDL-C test	80061, 83700, 83701, 83704, 83721	3048F, 3049F, 3050F
Schizophrenia	F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F25.0, F25.1, F25.8, F25.9	

Diabetes Monitoring for People with Diabetes and Schizophrenia (SMD)

The percentage of adults diagnosed with schizophrenia or schizoaffective disorder and diabetes who had both diabetes and cholesterol level tests during the measurement year. Measure population: Members ages 18–64 as of December 31 of the measurement year.

Required exclusions apply. See page 3.

Decription	СРТ	CPT-CAT-II
HbA1c tests	83036, 83037	3044F, 3046F, 3051F, 3052F
LDL-C tests	80061, 83700, 83701, 83704, 83721	3048F-3050F

Diabetes Screening for People with Schizophrenia or Bipolar Disorder who are Using Antipsychotic Medications (SSD)

The percentage of adults diagnosed with schizophrenia, schizoaffective disorder or bipolar disorder, who were dispensed an antipsychotic medication and had a test for diabetes during the measurement year. Measure population: Members ages 18–64 as of December 31 of the measurement year.

Decription	СРТ	CPT-CAT-II
HbA1c tests	83036, 83037	3044F, 3046F, 3051F, 3052F
Glucose tests	80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951	



Follow-Up After Emergency Department Visit for Substance Use Disorder or Dependence (FUA)

The percentage of ED visits for members 13 and older with a principal diagnosis of substance use disorder (SUD) or any diagnosis of drug overdose for which there was a follow-up visit.

Two rates are reported for follow-up visits after an ED visit:

- Within **7 days** of the ED visit (**8 total days**)
- Within 30 days of the ED visit (31 total days)

The follow-up visit may occur on the date of discharge and be with any practitioner with a principal diagnosis of SUD. If the first follow-up visit is within seven days after discharge, then both rates are counted for this measure. A pharmacotherapy dispensing event of alcohol use disorder treatment medications or opioid use disorder treatment medications within 7–30 days would also make the member compliant.

Description	СРТ	HCPCS	POS	ICD-10
Outpatient visit (with outpatient POS code) with any SUD diagnosis <u>or</u> Outpatient visit (with outpatient POS code) with a mental health provider	90847, 90853, 99238, 99239, 90875, 90876, 99223, 99222, 99221, 99255, 99253, 99254, 99252, 90849, 90791, 90792, 90845, 90840, 90839, 90832, 90833, 90834, 90836, 90837, 90838, 99233, 99232, 99231		03, 05, 07, 09, 11–20, 22, 33, 49, 50, 71, 72	See SUD Dx Codes
BH outpatient visit with a mental health provider <u>or</u> BH outpatient visit with any diagnosis of SUD	99483, 98961, 98962, 98960, 99345, 99342, 99344, 99341, 99350, 99348, 99349, 99347, 99510, 99385, 99386, 99387, 99384, 99382, 99381, 99383, 99494, 99492, 99245, 99243, 99244, 99242, 99205, 99203, 99204, 99202, 99211, 99215, 99213, 99214, 99212, 99395, 99396, 99397, 99394, 99392, 99391, 99393, 99078, 99401, 99402, 99403, 99404, 99411, 99412, 99493	G0176, H0040, H0039, H0004, H0002, T1015, H0037, H0036, H2015, H2016, H2010, H2000, H2011, G0463, H0034, H0031, H2013, H2017, H2018, G0512, G0155, H2014, G0409, H2019, H2020, G0177		See SUD Dx Codes
Telephone visit or telehealth visit with a mental health provider <u>or</u> Telephone or telehealth visit with any diagnosis of SUD	98966–98968, 99441–99443 90847, 90853, 99238, 99239, 90875, 90876, 99223, 99222, 99221, 99255, 99253, 99254, 99252, 90849, 90791, 90792, 90845, 90840, 90839, 90832, 90833, 90834, 90836, 90837, 90838, 99233, 99232, 99231			See SUD Dx Codes
E-visit or virtual check-in with a mental health provider <u>or</u> E-visit or virtual check-in with any diagnosis of SUD	98970, 98971, 98972, 99421, 99422, 99423, 99444, 99457, 99458, 98981, 98980	G2252, G2012, G2251, G0071, G2250, G2010		See SUD Dx Codes
Peer support services (with SUD diagnosis)		G0140, G0177, H0024, H0025, H0038–H0040, H0046, H2014, H2023, S9445, T1012, T1016		



Follow-Up After Emergency Department Visit for Substance Use Disorder or Dependence (FUA) (continued)

Description	СРТ	HCPCS	POS	ICD-10
Substance use disorder service or counseling/ surveillance	99408, 99409	G0396, G0397, G0443, H0001, H0005, H0007, H0015, H0016, H0022, H0047, H0050, H2035, H2036, T1006, T1012		Z71.41, Z71.51
BH screening or assessment for SUD or mental health disorders	99408, 99409	G0396, G0397, G0442, G2011, H0001, H0002, H0031, H0049		
Substance use service		H0006, H0028		
AOD medication treatment		G2067–G2070, G2072, G2073, H0020, H0033, J0570–J0573, J0577, J0578, J2315, Q9991– Q9992, S0109		
SUD diagnosis				F10.xx–16.xx, F18.xx–F19.xx, T40.xx–T43.xx, T51.xx



Follow-Up After Emergency Department Visit for Mental Illness (FUM)

The percentage of ED visits for members 6 years and older with a principal diagnosis of mental illness who received a follow-up visit with any provider.

Two rates are reported for follow-up visits after an ED visit:

- Within 7 days of the ED visit (8 total days)
- Within 30 days of the ED visit (31 total days)

If the first follow-up visit is within seven days after discharge, then both rates are counted for this measure.

Required exclusions apply. See page 3.

Mental Illness Diagnosis Codes

F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F21, F22, F23, F24, F25.0, F25.1, F25.8, F25.9, F28, F29, F30.10, F30.11, F30.12, F30.13, F30.2, F30.3, F30.4, F30.8, F30.9, F31.0, F31.10, F31.11, F31.12, F31.13, F31.2, F31.30, F31.31, F31.32, F31.4, F31.5, F31.60, F31.61, F31.62, F31.63, F31.64, F31.70, F31.71, F31.72, F31.73, F31.74, F31.75, F31.76, F31.77, F31.78, F31.81, F31.89, F31.9, F32.0, F32.1, F32.2, F32.3, F32.4, F32.5, F32.81, F32.89, F32.9, F32.A, F33.0, F33.1, F33.2, F33.3, F33.40, F33.41, F33.42, F33.8, F33.9, F34.0, F34.1, F34.81, F34.89, F34.9, F39, F40.00, F40.01, F40.02, F40.10, F40.11, F40.210, F40.218, F40.220, F40.228, F40.230, F40.231, F40.232, F40.233, F40.240, F40.241, F40.242, F40.243, F40.248, F40.290, F40.291, F40.298, F40.8, F40.9, F41.0, F41.1, F41.3, F41.8, F41.9, F42.2, F42.3, F42.4, F42.8, F42.9, F43.0, F43.10, F43.11, F43.12, F43.20, F43.21, F43.22, F43.23, F43.24, F43.25, F43.29, F43.8, F43.81, F43.89, F43.9, F44.89, F53.0, F53.1, F60.0, F60.1, F60.2, F60.3, F60.4, F60.5, F60.6, F60.7, F60.81, F60.89, F60.9, F63.0, F63.1, F63.2, F63.3, F63.81, F63.89, F63.9, F68.10, F68.11, F68.12, F68.13, F68.8, F68.A, F84.0, F84.2, F84.3, F84.5, F84.8, F84.9, F90.0, F90.1, F90.2, F90.8, F90.9, F91.0, F91.1, F91.2, F91.3, F91.8, F91.9, F93.0, F93.8, F93.9, F94.0, F94.1, F94.2, F94.8, F94.9

Self-Harm Diagnosis Codes

X71-X83, T36-T65, T71, R45.851

CountyCare A MEDICAID HEALTH PLAN

Follow-Up After Emergency Department Visit for Mental Illness (FUM) (continued)

Description	СРТ	HCPCS	POS	ICD-10
Outpatient visit (with outpatient POS Value Set) with any diagnosis of mental health disorder	90847, 90853, 99238, 99239, 90875, 90876, 99223, 99222, 99221, 99255, 99253, 99254, 99252, 90849, 90791, 90792, 90845, 90840, 90839, 90832, 90833, 90834, 90836, 90837, 90838, 99233, 99232, 99231		See Mental Illness Dx codes	03, 05, 07, 09, 11–20, 22, 33, 49, 50, 71, 72
Behavioral health outpatient visit with any diagnosis of mental health disorder	99483, 98961, 98962, 98960, 99345, 99342, 99344, 99341, 99350, 99348, 99349, 99347, 99510, 99385, 99386, 99387, 99384, 99382, 99381, 99383, 99494, 99492, 99245, 99243, 99244, 99242, 99205, 99203, 99204, 99202, 99211, 99215, 99213, 99214, 99212, 99395, 99396, 99397, 99394, 99392, 99391, 99393, 99078, 99401, 99402, 99403, 99404, 99411, 99412, 99493	G0176, H0040, H0039, H0004, H0002, T1015, H0037, H0036, H2015, H2016, H2010, H2000, H2011, G0463, H0034, H0031, H2013, H2017, H2018, G0512, G0155, H2014, G0409, H2019, H2020, G0177	See Mental Illness Dx codes	
Telehealth or telephone visit with any diagnosis of mental health disorder	98966–98968, 99441–99443 90847, 90853, 99238, 99239, 90875, 90876, 99223, 99222, 99221, 99255, 99253, 99254, 99252, 90849, 90791, 90792, 90845, 90840, 90839, 90832, 90833, 90834, 90836, 90837, 90838, 99233, 99232, 99231		See Mental Illness Dx codes	02
Transitional care management services with any diagnosis of mental health disorder	99495, 99496		See Mental Illness Dx codes	
Psychiatric collaborative care management	99492, 99493, 99494	G0512		
E-visit or virtual check-in with any diagnosis of mental health disorder	98970, 98971, 98972, 99421, 99422, 99423, 99444, 99457, 99458, 98981, 98980	G2252, G2012, G2251, G0071, G2250, G2010	See Mental Illness Dx codes	



Substance Use Disorder (FUI)

This measure looks at the percentage of acute inpatient hospitalizations, residential treatment or detoxification visits for a diagnosis of substance use disorder among members 13 years of age and older that result in a follow-up visit or service for substance use disorder.

Two rates are reported:

- The percentage of visits or discharges for which the member received follow-up for substance use disorder within the **30 days** after the visit or discharge.
- The percentage of visits or discharges for which the member received follow-up for substance use disorder within the **7 days** after the visit or discharge.

Required exclusions apply: Members in hospice are excluded from the eligible population.

The follow-up visit or event may be with any practitioner for a principal diagnosis of substance use disorder. Do not include visits that occur on the date of discharge.

Report three age stratifications and a total rate:

- 13–17 years
- 18 64 years
- 65 years and older
- Total

Outpatient Follow-Up Visit

CPT: 90847, 90853, 99238, 99239, 90875, 90876, 99223, 99222, 99221, 99255, 99253, 99254, 99252, 90849, 90791, 90792, 90845, 90840, 90839, 90832, 90833, 90834, 90836, 90837, 90838, 99233, 99232, 99231

Outpatient POS

03, 05, 07, 09, 11-20, 22, 33, 49, 50, 71, 72

BH Outpatient Visit

CPT: 99483, 98961, 98962, 98960, 99345, 99342, 99344, 99341, 99350, 99392, 99391, 99393, 99078, 99401, 99402, 99403, 99404, 99411, 99383, 99494, 99492, 99245, 99243, 99244, 99242, 99205, 99203, 99412, 99493, 99204, 99348, 99349, 99347, 99510, 99385, 99386, 99387, 99384, 99382, 99381, 99202, 99211, 99215, 99213, 99214, 99212, 99395, 99396, 99397, 99394

Telephone Visit

CPT: 98966, 98967, 98968, 99441, 99442, 99443

Online Assessment

CPT: 98969, 98970, 98971, 98972, 99421, 99422, 99423, 99444, 99457, 99458

HCPCS: G0071, G2010, G2012, G2250, G2251, G2252

Substance Use Disorder Diagnosis

ICD-10: F10.xx-16.xx, F18.xx-19.xx

CountyCare A MEDICAID HEALTH PLAN

Follow-Up After Hospitalization for Mental Illness (FUH)

The percentage of discharges for members 6 years and older who were hospitalized due to a principal diagnosis of mental illness or any diagnosis of self-harm and who had a timely mental health follow-up visit.

Two rates are reported for:

- Follow-up care within **7 days** after discharge.
- Follow-up care within 30 days after discharge.

Do not include visits that occur on the date of discharge.

Mental Illness Diagnosis

F20.0, F20.1, F20.2, F20.3, F20.5, F20.81, F20.89, F20.9, F21, F22, F23, F24, F25.0, F25.1, F25.8, F25.9, F28, F29, F30.10, F30.11, F30.12, F30.13, F30.2, F30.3, F30.4, F30.8, F30.9, F31.0, F31.10, F31.11, F31.12, F31.13, F31.2, F31.30, F31.31, F31.32, F31.4, F31.5, F31.60, F31.61, F31.62, F31.63, F31.64, F31.70, F31.71, F31.72, F31.73, F31.74, F31.75, F31.76, F31.77, F31.78, F31.81, F31.89, F31.9, F32.0, F32.1, F32.2, F32.3, F32.4, F32.5, F32.81, F32.89, F32.9, F32.A, F33.0, F33.1, F33.2, F33.3, F33.40, F33.41, F33.42, F33.8, F33.9, F34.0, F34.1, F34.81, F34.89, F34.9, F39, F40.00, F40.01, F40.02, F40.10, F40.11, F40.210, F40.218, F40.220, F40.228, F40.230, F40.231, F40.232, F40.233, F40.240, F40.241, F40.242, F40.243, F40.248, F40.290, F40.291, F40.298, F40.8, F40.9, F41.0, F41.1, F41.3, F41.8, F41.9, F42.2, F42.3, F42.4, F42.8, F42.9, F43.0, F43.10, F43.11, F43.12, F43.20, F43.21, F43.22, F43.23, F43.24, F43.25, F43.29, F43.8, F43.81, F43.89, F43.9, F44.89, F53.0, F53.1, F60.0, F60.1, F60.2, F60.3, F60.4, F60.5, F60.6, F60.7, F60.81, F60.89, F60.9, F63.0, F63.1, F63.2, F63.3, F63.81, F63.89, F63.9, F68.10, F68.11, F68.12, F68.13, F68.8, F68.A, F84.0, F84.2, F84.3, F84.5, F84.8, F84.9, F90.0, F90.1, F90.2, F90.8, F90.9, F91.0, F91.1, F91.2, F91.3, F91.8, F91.9, F93.0, F93.8, F93.9, F94.0, F94.1, F94.2, F94.8, F94.9

Self Harm

X71-X83, T36-T65, T71, R45.851

Description	СРТ	HCPCS	ICD-10	POS
Outpatient visit (with outpatient POS value set) with a mental health provider <u>or</u> Outpatient visit (with outpatient POS value set) with any diagnosis of mental health disorder	90847, 90853, 99238, 99239, 90875, 90876, 99223, 99222, 99221, 99255, 99253, 99254, 99252, 90849, 90791, 90792, 90845, 90840, 90839, 90832, 90833, 90834, 90836, 90837, 90838, 99233, 99232, 99231		See Mental Illness Dx codes	03, 05, 07, 09, 11–20, 22, 33, 49, 50, 71, 72
Behavioral health outpatient visit with a mental health provider <u>or</u> Behavioral health outpatient visit with any diagnosis of mental health disorder	99483, 98961, 98962, 98960, 99345, 99342, 99344, 99341, 99350, 99348, 99349, 99347, 99510, 99385, 99386, 99387, 99384, 99382, 99381, 99383, 99494, 99492, 99245, 99243, 99244, 99242, 99205, 99203, 99204, 99202, 99211, 99215, 99213, 99214, 99212, 99395, 99396, 99397, 99394, 99392, 99391, 99393, 99078, 99401, 99402, 99403, 99404, 99411, 99412, 99493	G0176, H0040, H0039, H0004, H0002, T1015, H0037, H0036, H2015, H2016, H2010, H2000, H2011, G0463, H0034, H0031, H2013, H2017, H2018, G0512, G0155, H2014, G0409, H2019, H2020, G0177	See Mental Illness Dx codes	



Follow-Up After Hospitalization for Mental Illness (FUH) (continued)

Description	СРТ	HCPCS	ICD-10	POS
Telehealth or telephone visit with a mental health provider (use POS value for telehealth) <u>or</u> A telehealth or telephone visit with any diagnosis of mental health disorder	98966–98968, 99441–99443 90847, 90853, 99238, 99239, 90875, 90876, 99223, 99222, 99221, 99255, 99253, 99254, 99252, 90849, 90791, 90792, 90845, 90840, 90839, 90832, 90833, 90834, 90836, 90837, 90838, 99233, 99232, 99231		See Mental Illness Dx codes	02
Transitional care management services with a mental health disorder <u>or</u> Transitional care management services with any diagnosis of mental health disorder	99495, 99496		See Mental Illness Dx codes	
Psychiatric collaborative care management	99492, 99493, 99494	G0512		

Providers that qualify as Mental Health Provider include:

- An MD, DO, APN, or PA who specializes in Psychiatry for children or Adults.
- An RN who is certified and credentialed as a Psychiatric Nurse or Mental Health Clinical Nurse Specialist.
- A Licensed Psychologist, Therapist, Counselor (Including LPCs & LCPCs), or Social Worker (LCSW).
- A Certified Community Behavioral Health Center or Certified Community Behavioral Health Clinic.



Follow-Up Care for Children Prescribed ADHD Medication (ADD-E)

The percentage of children newly prescribed with attention deficit/hyperactivity disorder (ADHD) medication who received follow-up care. Two rates are reported.

- Initiation Phase. The percentage of members with an outpatient prescription dispensed for ADHD medication, who had one follow-up visit with a prescribing practitioner within **30 days** following the IPSD.
- Continuation and Maintenance (C&M) Phase. The percentage of members with an outpatient prescription dispensed for ADHD medication, who remained on the medication for 210 days or more, and who had two additional follow-up visits with a practitioner within 270 days after the end of the Initiation Phase.

Measure population: Members ages 6–12 as of the Index Prescription Start Date (IPSD), the earliest ADHD medication dispense date.

Required exclusions apply: See page 3.

Description СРТ **HCPCS** POS **Outpatient visit (Visit Setting** 90791, 90792, 90832-90834, 90836-03, 05, 07, 90840, 90845, 90847, 90849, 90853, 09, 11–20, Unspecified Value Set with **Outpatient POS Value Set)** 90875, 90876, 99221-99223, 99231-22, 33, 49, 99233, 99238, 99239, 99251–99255 50, 71, 72 **Outpatient visit (Behavioral Health** 98960-98962, 99078, 99202-99205, G0155, G0176, G0177, 99211-99215, 99241-99245, 99341-Outpatient Value Set) G0409, G0463, H0002, 99345, 99347-99350, 99381-99387, H0004, H0031, H0034, 99391-99397, 99401-99404, 99411, H0036, H0037, H0039, 99412, 99510, 99483 H0040, H2000, H2010, H2011, H2013-H2020, T1015 Telehealth or telephone visits 98966-98968, 99441-99443 02 (use POS value for telehealth)

Can be in-person, Telephone or Telehealth visit.



Initiation & Engagement of Alcohol and Other Drug Abuse or Independence Treatment (IET)

The percentage of adolescent and adult members with a new episode of alcohol or other drug (AOD) abuse or dependence who received the following:

- Initiation of AOD Treatment. Initiated dependence treatment within 14 days of their diagnosis.
- Engagement of AOD Treatment. Continued treatment with two or more additional services within 34 days of the initiation visit.

Measure population: Medicaid and Medicare members ages 13 or older as of December 31 of the measurement year. For the follow-up treatments, include an ICD-10 diagnosis for alcohol or other drug dependence from the Mental, Behavioral and Neurodevelopmental Disorder Section of ICD-10 along with a procedure code for the preventive service, evaluation and management consultation or counseling service.

Exclusions: Members in hospice are excluded from the eligible population.

Description	СРТ	HCPCS	POS	ICD-10
Telephone visit and Telehealth Visit	98966–98968, 99441–99443		02	
Online assessment	98969–98972, 99421–99423, 99444, 99458	G2010, G2012		
Alcohol and other drug medication treatment	98970–98972, 99421, 99422, 99423, 99458	H0020, H0033, J0570, J0571–J0575, J2315, Q9991, Q9992, S0109		
Substance use disorder diagnosis				F10.xx–16.xx, F18.xx–19.xx

Can be in-person or Telephone visit.

Metabolic Monitoring for Children and Adolescents on Antipsychotics (APM)

The percentage of children and adolescents who had two or more antipsychotic prescriptions and had metabolic testing.

Measure population: Medicaid members ages 1–17 as of December 31 of the measurement year.

Three rates are reported:

- Blood glucose or HbA1c testing
- Cholesterol or LDL-C testing
- Blood glucose and cholesterol testing

Description	СРТ	CPT-CAT-II
HbA1c	83036, 83037	3044F, 3046F, 3051F, 3052F
Glucose	80047, 80048, 80050, 80053, 80069, 82947, 82950, 82951	
LDL-C	80061, 83700, 83701, 83704, 83721	3048F, 3049F, 3050F



Pharmacotherapy for Opioid Use Disorder (POD)

This measure captures the percentage of new opioid use disorder (OUD) pharmacotherapy events with OUD pharmacotherapy for **180 or more days** among members age 16 and older with a diagnosis of OUD.

- The OUD dispensing event will be captured between a 12-month period that begins on July 1 of the year prior to the measurement year and ends on June 30 of the measurement year (Intake Period). This ensures capture of pharmacotherapy compliance by December 31 of the measurement year.
- Members must have a Negative Medication History (no OUD pharmacotherapy medications captured on pharmacy claims) as of **31 days** prior to the new OUD pharmacotherapy to be included in the measure population.

Measure population: Members 6 years and older as of December 31 of the measurement year. Report two age stratifications and total rate:

- 16-64 years
- 65 years and older
- Total

Required exclusions apply. See page 3.

Medical Record Documentation and Best Practices:

Build a partnership on trust and understanding with the patient.

- Medication regiment adherence is essential for the patient's treatment.
- Provide credible sources in order to address any fears and stigma surrounding treatment.
- Recognize that the patient might want to participate at varying levels, so meet them where they are.
- Decision making should include the patient and their family.

Description	Prescription	Value Sets and Days Supply
Antagonist	Naltrexone (oral)	NA—Codes do not exist
Antagonist	Naltrexone (injectable)	Naltrexone Injection Value Set (31 days supply)
Partial agonist	Buprenorphine (sublingual tablet)	Buprenorphine Oral Value Set (1 day supply) Buprenorphine Oral Weekly Value Set (7 days supply)
Partial agonist	Buprenorphine (injection)	Buprenorphine Injection Value Set (31 days supply)
Partial agonist	Buprenorphine (implant)	Buprenorphine Implant Value Set (180 days supply)
Partial agonist	Buprenorphine/naloxone (sublingual tablet, buccal film, sublingual film)	Buprenorphine Naloxone Value Set (1 day supply)
Agonist	Methadone (oral)	Methadone Oral Value Set (1 day supply) Methadone Oral Weekly Value Set (7 days supply)

Methadone is not included on the medication lists for this measure. Methadone for OUD administered or dispensed by federally certified opioid treatment programs (OTP) is billed on a medical claim. A pharmacy claim for methadone would be indicative of treatment for pain rather than OUD.



